

Syddansk Universitet

Identification of PPAR agonists in root extracts of purple coneflower (*Echinacea purpurea*) by bioassay-guided fractionation

El-Houri, Rime Bahij; Fretté, Xavier; Christensen, Kathrine Bisgaard; Kotowska, Dorota Ewa; Grevsen, Kai; Kristiansen, Karsten; Christensen, Lars Porskjær

Publication date:
2011

Document Version
Final published version

[Link to publication](#)

Citation for pulished version (APA):

El-Houri, R. B., Fretté, X., Christensen, K. B., Kotowska, D. E., Grevsen, K., Kristiansen, K., & Christensen, L. P. (2011). Identification of PPAR agonists in root extracts of purple coneflower (*Echinacea purpurea*) by bioassay-guided fractionation. Abstract from Nordic Natural Products Conference (NNPC) , Holmen, Oslo, Norway.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Identification of PPAR γ agonists in root extracts of purple coneflower (*Echinacea purpurea*) by bioassay-guided fractionation

Rime Bahij El-Houri^{1*}, X. C. Fretté¹, K. B. Christensen¹, D. Kotowska², K. Grevsen³, K. Kristiansen² and L. P. Christensen¹

Introduction

One of the major characteristics of type 2 diabetes (T2D) is insulin resistance, which is often treated by insulin sensitizing drugs such as thiazolidinediones (TZDs). The primary target for the TZDs is the peroxisome proliferator-activated receptor PPAR γ . However, critical side effects of TZDs can occur, as they are full PPAR γ agonists. Partial PPAR γ agonists are associated with fewer side effects but still may maintain the effect on insulin resistance. Alkamides are very similar in chemical structure to natural ligands for the PPAR γ such as fatty acids, and hence are potential PPAR γ agonists. Recently, a new C₁₆-alkamide able to activate PPAR γ with no concurrent stimulation of adipocyte differentiation, and able to increase insulin-stimulated glucose uptake was isolated from the flowers of purple coneflower (*Echinacea purpurea*) [1]. In our search for new partial PPAR γ agonists the roots of purple coneflower, which is a rich source for alkamides, was investigated.



Extraction and isolation

5 kg fresh roots of purple coneflower were extracted overnight with dichloromethane (DCM). The dried extract was subjected to bioassay-guided chromatographic fractionation. First step was a separation by flash column chromatography using a *n*-hexane–ethyl acetate gradient resulting in 10 fractions (A–J). Two fractions (D and H) showed promising bioactivity in a dose-dependent manner. LC-MS analysis revealed that these two fractions primarily contained known and unknown alkamides (data not shown).




PPAR γ transactivation assay

The ability of the DCM extract to activate the PPAR γ was tested in a luciferase-based PPAR transactivation assay using mouse embryonic fibroblasts transfected with a luciferase reporter plasmid, a transfection control plasmid, and an expression plasmid. Degree of activation was determined by a luminometer and compared to a positive control, Rosiglitazone (Rosi). The DCM extract was found to activate PPAR γ in a dose-dependent manner (1–100 μ g/mL) without stimulating adipocyte differentiation at 100 μ g/mL. Figures 1 and 2 show fold activations of PPAR γ by the DCM extract with vehicle (DMSO) set to 1.0 and corresponding results from adipocyte differentiation assay (DEX protocol).



Stimulation of glucose uptake in adipocytes

The DCM extract and fractions D and H were tested for stimulation of insulin-dependent glucose uptake in adipocytes. Dose-dependent stimulation (0–100 nM) was observed for all three samples as shown in Figure 3. All results were compared to a positive control (Rosi) and vehicle (DMSO). Mature adipocytes were subjected to the extract/fractions and after 2 days glucose uptake was induced by insulin. Effect was measured using ¹⁴C-labeled glucose and afterwards radioactivity was determined by scintillation counting.



Conclusions and Perspectives

- The DCM extract of the roots of purple coneflower showed promising activation of PPAR γ in a dose-dependent manner without stimulating adipocyte differentiation.
- Bioassay-guided fractionation resulted in two alkamide-rich fractions, which were able to activate PPAR γ and positively affected insulin-stimulated glucose uptake in adipocytes.
- The results of this study clearly indicate that alkamides are partial PPAR γ agonists with a potential in relation to the management of insulin resistance and T2D. Individual alkamides from the active fractions warrant further investigation for their PPAR γ activating properties in order to identify the most active alkamides and hence, the most promising PPAR γ agonists in coneflower roots.

Reference

[1] Christensen KB et al. *J. Nat. Prod.* 2009; 72: 933–937.

¹Institute of Chemical Engineering, Biotechnology & Environmental Technology, University of Southern Denmark, Niels Bohrs Allé 1, DK-5230 Odense M. E-mail: rbeh@kbm.sdu.dk. ²Dept. of Biology, Copenhagen University, ³Dept. of Horticulture, Aarhus University

